

References

Nayeli Macias, Heliodoro Alemán-Mateo, Julián Esparza-Romero and Mauro E Valencia (2007). Body fat measurement by bioelectrical impedance and air displacement plethysmography: a cross-validation study to design bioelectrical impedance equations in Mexican adults. *Nutrition Journal*, 6:18:10.1186/1475-2891-6-18.

Klauke S, Fischer H, Rieger A, Frühauf L, Staszewski S, Althoff PH, Helm EB (2005). Use of bioelectrical impedance analysis to determine body composition changes in HIV-associated wasting. *Int J STD AIDS*. Apr;16(4):307-13.

Sun SS, Chumlea WC, Heymsfield SB, Lukaski HC, Schoeller D, Friedl K, Kuczmarski RJ, Flegal KM, Johnson CL, Hubbard VS (2003). Development of bioelectrical impedance analysis prediction equations for body composition with the use of a multicomponent model for use in epidemiologic surveys. *Am J Clin Nutr*. Feb;77(2):331-40.

ADAMS, J.A. (1985) Ross Medical Publications. Symposium on human body composition.

COHN, D.A., KAY, T.P., TATSCH, R.F., THIES, C.F., (1981). Comparison of methods for estimating body fat in normal subjects and cancer patients. *AMERICAN JOURNAL OF CLINICAL NUTRITION*, vol. 34, 2839-2847.

DURIN, J.U.G.A., WORMERSLY, J., (1974). Body fat assessment from total body density and its estimation from skinfold thickness: Measurement of 481 men and women aged 16-72. *BRITISH JOURNAL OF NUTRITION*, vol 32, 77-97.

GIRANDOLA, R.N., CONTARSI, S.A., WISWELL, R.A., ARTAL, R. MD, Algorithmic ElectroLipoGraphy. A simple and accurate method for the clinical diagnosis of obesity.

HARSHA, D.W., VOORS, A.W., BERENSON, G.S., (1978). Racial differences in subcutaneous fat patterns in children age 7-15. *AMERICAN JOURNAL OF PHYSICAL ANTHROPOLOGY*, VOL 53, 333-337.

HARSHA, D.W., FREDRICKS, R., BENSON, G.S., (1978). Denisometry and anthropometry of blacks and whites. *Human Biology*, vol. 50, 261-280.

KATCH, F.I., KATCH, V.L., (1980). Measurement and prediction errors in body composition assessment and the search for the perfect equation. RESEARCH QUARTERLY, vol. 51, no. 1, 249-260.

THOMASSETT, M.A., (1963). Properties of Bio-Electrical Impedance des tissue. LYON MEDICINE, VOL. 22, 11325-1350.

LUKASKI, H.C., MENDEZ, J., Assessment of fat-free mass using Bio-Electrical Impedance measurements of the human body. AMERICAN

JOURNAL OF CLINICAL NUTRITION, vol. 41, no. 4, 809-816.

SEGAL, K.R., GUTIN, B., PRESTA, E., WANG, J., VAN ITALLIE, T.B., (1988). Estimation of human body composition by electrical impedance methods: a comparative study. St. Lukes-Roosevelt Hospital Center and Applied Physiology Lab.

JACKSON, A.S., POLLOCK, M.L. GRAVES, J.E., MAHAR, M.T., (1988). Reliability and validity of bioelectrical impedance in determining body composition. Department of Health and Physical Education, and Recreation, University of Houston.

ISRAEL, R.G., HOUMARD, J.A., O'BRIENS, K.F. (1990). Validity of NIR for estimating human body composition. MEDICINE AND SCIENCE IN SPORTS AND EXERCISE, vol 21, no.2, S103.

DAVIS, P.G., VAN LOAN, M., HOLLY, R.G., KRSTICH, K., PHINNEY, S.D., (1990). Near infrared interactance VS. hydrostatic weighing to measure body composition in lean, normal and obese women. MEDICINE AND SCIENCE IN SPORTS AND EXERCISE, vol. 21, no.2, S13.